

REQUEST FOR ACCESS TO ABANDONED AP. 20

UNDER 37 CFR 1.14(a)

~~RECEIVED~~

DEC 06 1999

File Information Unit

Unit 1B Application of

ADMISSION NUMBER	07/310252	FILED 2/13/89
GATE AT LINE	Examiner 184	Feisee 381

Assistant Commissioner for Patents
Washington, DC 20231

I hereby request access under 37 CFR 1.14(a)(3)(vi) to the application file record of the above-identified ABANDONED application, which is: (CHECK ONE) **5530101** column _____

(A) referred to in United States Patent Number

5530/01

scilicet

(A) referred to in United States Patent Number _____
 (B) referred to in an application that is open to public inspection as set forth in 37 CFR 1.11, i.e., Application No. _____, filed _____, on page _____ of paper number _____.
 (C) an application that claims the benefit of the filing date of an application that is open to public inspection, i.e., Application No. _____, filed _____.
 (D) an application in which the applicant has filed an authorization to lay open the complete application to the public.
following address:

Please direct any correspondence concerning this request to the following address:

Please direct any correspondence concerning this request to the following:

Ramón J. Ríos
Signature

ສິນຄະນະ

RICHARD AREIA

Types of sorted name

12-6-99

8818

~~FOR STO USE ONLY~~

Approved by

Onassis

Unit 2

United States Patent [19]
Queen et al.

US005530101A

[11] Patent Number: 5,530,101
[45] Date of Patent: Jun. 25, 1996

[54] HUMANIZED IMMUNOGLOBULINS

[75] Inventors: Cary L. Queen, Los Altos; Harold E. Selick, Belmont, both of Calif.

[73] Assignee: Protein Design Labs, Inc., Mountain View, Calif.

[21] Appl. No.: 634,278

[22] Filed: Dec. 19, 1990

Related U.S. Application Data

[63] Continuation-in-part of Ser. No. 590,274, Sep. 28, 1990, abandoned, and a continuation-in-part of Ser. No. 310,252, Feb. 13, 1989, abandoned, which is a continuation-in-part of Ser. No. 290,975, Dec. 28, 1988, abandoned.

[51] Int. Cl. 461K 39/395; C07K 16/28

[52] U.S. Cl. 530/387.3; 530/387.1;
530/388.22; 424/133.1; 424/143.1

[58] Field of Search 424/85.8, 133.1,
424/143.1; 530/387, 388.22, 387.1, 387.3

[56] References Cited

U.S. PATENT DOCUMENTS

4,816,397 3/1989 Boss et al. 435/68
4,816,567 3/1989 Cabilly et al. 530/387
4,867,973 9/1989 Geers et al.
5,225,539 7/1993 Winter .

FOREIGN PATENT DOCUMENTS

0171496 2/1986 European Pat. Off.
0173494 3/1986 European Pat. Off.
0184187 6/1986 European Pat. Off.
0239400 9/1987 European Pat. Off.
0266663 6/1988 European Pat. Off.
2188941 10/1987 United Kingdom .
WO86/05513 9/1986 WIPO .
WO87/02671 5/1987 WIPO .
WO89/01783 3/1989 WIPO .

OTHER PUBLICATIONS

Vitteta et al., "Redesigning Nature's Poisons to Create Anti-Tumor Reagents," *Science* 238:1098-1104 (1987). Ellison et al., "The nucleotide sequence of a human immunoglobulin C(gamma)₁ gene", *Nucleic Acids Res.* 10:4071-(1982).

Hicter et al., "Cloned Human and Mouse Kappa Immunoglobulin Constant and J Region Genes Conserves homology in Functional Segments", *Cell* 22:197-207 (1980).

Sharon et al., "Expression of a V_HC_K chimaeric protein in mouse myeloma cells", *Nature* 309:364-367 (1984).

Takeda et al., "Construction of chimaeric processed immunoglobulin genes containing mouse variable and human constant region sequences", *Nature* 314:452-454 (1985).

Tan et al., "A Human-Mouse Chimeric Immunoglobulin Gene with a Human Variable Region is Expressed in Mouse Myeloma Cells", *J. Immunol.* 135:3564-3567 (1985).

Morrison et al., "Chimeric human antibody molecules: Mouse antigen-binding domains with human constant region domains," *Proc. Natl. Acad. Sci.* 81:6851-6859 (1984).

Boulianne et al., "Production of functional chimeric mouse/human antibody," *Nature* 312:643-646 (1984).

Neuberger et al., "A haptene-specific chimeric IgE antibody with human physiological effector function," *Nature* 314:268-270 (1985).

Morrison, S. L., "Transfectomas Provide Novel Chimeric Antibodies," *Science* 229:1202-1207 (1985).

Sahagian et al., "A Genetically Engineered Murine/Human Chimeric Antibody Retains Specificity for Human Tumor-Associated Antigen", *J. Immunol.* 137:1066-1074 (1986). Liu et al., "Expression of mouse:human immunoglobulin heavy-chain cDNA in lymphoid cells", *Gene* 54:33-40 (1987).

Better et al., "Escherichia coli Secretion of an Active Chimeric Antibody Fragment", *Science* 240:1041-1043 (1988).

Waldmann, T. A., "The Structure, Function, and Expression of Interleukin-2 Receptors on Normal and Malignant Lymphocytes," *Science* 232:727-732 (1986).

Leonard et al., "The human receptor for T-cell growth factor," *J. Biol. Chem.* 260:1872-1880 (1985).

Farrar, J., "The biochemistry, biology, and role of interleukin-2 in the induction of cytotoxic T cell and antibody-forming B cell receptors," *Immunol. Rev.* 63:129-166 (1982).

Greene et al., "Growth of Human T Lymphocytes: An Analysis of Interleukin 2 and Its Cellular receptor", in *Progress in Hematology XIV*, E. Brown ed., Grune and Stratton, New York (1986) pp. 283-301.

Verhoyen et al., "Reshaping Human Antibodies: Grafting an Antilysozyme Activity", *Science* 239:1534-1536 (1988).

Joncs et al., "Replacing the complementarity-determining regions in a human antibody with those from a mouse", *Nature* 321:522-525 (1986).

Hale et al., "Remission Induction in Non-Hodgkin Lymphoma with Reshaped Human Monoclonal Antibody CAMPATH-1H", *Lancet* Dec. 17, 1988, pp. 1394-1399.

Chothia, C. and A. M. Lesk, "Canonical Structures for the Hypervariable Regions of Immunoglobulins", *J. Mol. Biol.* 196:901-917 (1987).

(List continued on next page.)

Primary Examiner—Lila Fcisic
Attorney, Agent, or Firm—Townsend and Townsend and Crew

[57]

ABSTRACT

Novel methods for producing, and compositions of, humanized immunoglobulins having one or more complementarity-determining regions (CDR's) and possible additional amino acids from a donor immunoglobulin and a framework region from an accepting human immunoglobulin are provided. Each humanized immunoglobulin chain will usually comprise, in addition to the CDR's, amino acids from the donor immunoglobulin framework that are, e.g., capable of interacting with the CDR's to effect binding affinity, such as one or more amino acids which are immediately adjacent to a CDR in the donor immunoglobulin or those within about 3 Å as predicted by molecular modeling. The heavy and light chains may each be designed by using any one or all of various position criteria. When combined into an intact antibody, the humanized immunoglobulins of the present invention will be substantially non-immunogenic in humans and retain substantially the same affinity as the donor immunoglobulin to the antigen, such as a protein or other compound containing an epitope.